

Orange molasses as a new energy ingredient for feedlot lambs in Brazil

ISABELA JORGE SANTOS (✉ isabelajorge@usp.br)

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz
<https://orcid.org/0000-0002-2772-1495>

Paulo César Gonzales Dias Junior

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

Ana Carolina Silva Vicente

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

Adrielly Lais Alves

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

Rhaissa Garcia de Assis

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

Janaina Socolovski Biava

ESALQ-USP: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

Mateus Vigo Vercesi Almada Nogueira

cpKELCO

Alexandre Vaz Pires

USP FMVZ: Universidade de Sao Paulo Faculdade de Medicina Veterinaria e Zootecnia

Evandro Maia Ferreira

USP ESALQ: Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz
<https://orcid.org/0000-0002-1057-5508>

Research Article

Keywords: Sheep, Performance, By-product, Replacement, Flint corn, Feed efficiency

Posted Date: March 6th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2499632/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Tropical Animal Health and Production on July 3rd, 2023. See the published version at <https://doi.org/10.1007/s11250-023-03675-4>.

Abstract

The objectives of this experiment were to evaluate the effects of increasing levels of orange molasses in replacement of flint corn grain in high-concentrate diets on dry matter intake (DMI), average daily gain (ADG) and feed efficiency (FE) of feedlot lambs. Thirty male lambs without defined racial pattern (30.3 ± 5.3 kg of initial BW; mean \pm SD) were used in a randomized complete block design with 10 blocks and 3 treatments. The treatments were defined by partial replacement of flint corn by orange molasses in the diet with 90% of concentrate and 10% of *Cynodon spp* hay, as follow: **00M** – control diet without orange molasses; **200M** – 20% of orange molasses replacing flint corn and **400M** – 40% of orange molasses replacing flint corn (DM basis). The experiment lasted 72 days divided into 3 subperiods, with 1 subperiod of 16 days and 2 subperiods of 28 days. Animals were weighed after a 16-h fast on days 1, 16, 44 and 72 of the experimental periods to determine the ADG and FE. The DMI, ADG and FE showed an interaction between treatments and experimental periods. The DMI in the first period decreased linearly ($P < 0.01$), in the third period, there was no effect of treatments ($P > 0.05$) on DMI. The ADG decreased linearly ($P < 0.01$) in the first period as the orange molasses increased. Otherwise, in the third period, ADG increased linearly ($P = 0.05$) as flint corn was replacement by orange molasses. The FE showed an interaction between treatment and period ($P = 0.09$). In the first period had a decreased linear effect, in the third period there was a trend ($P = 0.07$) of linear increased. There was no difference between the diets regarding the final BW of the lambs. In conclusion, the orange molasses can replace up to 40% of flint corn in diets for feedlot lambs without affecting final BW. However, it is important to consider the adaptation time proved to be very important for better use of orange molasses as a source of energy in diets for lambs.

1. Introduction

Finish lambs on feedlot receiving a high-concentrate diets are a technique that increases the ADG and reduces the slaughter time. However, these diets are expensive. Corn grain is the main source of energy used in ruminant diets in Brazil, 97% of feedlot nutritionists use corn as source of energy (Silvestre and Millen, 2021). Corn is the main agricultural commodity that impacts the cost of farm animal diets. According to USDA (United States Department of Agriculture), corn consumption in the 2019/2020 harvest reached 1.127 billion tons higher than total production, resulting in a reduction in world stock. An alternative to reduce the cost of animal feed and dependence on corn is the use of agro-industry by-products (De Evan et al., 2020; Vastolo et al., 2022). The orange molasses is a by-product of the citrus processing industry, which produces many by-products that are already used in animal feed, having citrus pulp as the most commonly used by-product for ruminants (Fegeros et al., 1995; Arthington et al., 2002; Santos et al., 2014).

Brazil is the world's largest orange juice producer, accounting for more than one quarter of the world production (Pires et al., 2021). In this context, it produces a large number of co-products, including orange molasses. The Orange molasses is made by concentrating the press liquor from the citrus peel residue. It is an ingredient with high moisture content, which has about 100 – 170 g/kg solubles of which 500 –

700 g/kg consists of sugar (Ensminger et al., 1990). The orange molasses is a safe food with high nutritional potential and can be a source of energy in the ruminant diet, but it is not yet explored for animal production. It is mainly destined for the food industry as an emulsifying agent for pellets and for fertigation (Tuttobene et al., 2009). There are studies in the literature that evaluated the effects of orange molasses in the diet of cows, steers, and lambs (Becker et al., 1944; Chen et al., 1981; Wing et al., 1988). However, there is a lack of information on the effects of orange molasses in high-concentrate diets for lambs.

The use of orange molasses in diets for ruminants can reduce dependence on corn, the main ingredient in finishing diets and whose price is currently high. Furthermore, this article has an environmental appeal, as the use of by-products from the industry in animal feed contributes to the lowest disposal in the environment. Thus, this study evaluated the partial replacement of flint corn by orange molasses on the performance of feedlot lambs.

2. Material And Methods

This study was carried out at the at the sheep confinement facility of the Sheep and Goat Intensive Production System (SIPOC) of Animal Science Department “Luiz de Queiroz” College of Agriculture, University of São Paulo (ESALQ-USP), located in Piracicaba, SP, Brazil, at altitude of 540 m and average annual temperature of 24.40 °C. The research protocol was approved by the Animal Care and Use Committee (number 7366010422).

Thirty male lambs with no defined racial pattern and BW 30.3 ± 3.07 kg (mean \pm SD) were used in a randomized completely block design (10 blocks and 3 treatments). The control diet contained 70.6% flint corn and 0% orange molasses (**00M**). In the remaining diets, orange molasses replaced flint corn at the rate of 20 (**200M**) or 40% (**400M**) of the original corn concentration, resulting in 0, 14.1, or 28.2% of orange molasses in the dietary DM (Table 1). The experimental diets were formulated to be isonitrogenous (Table 1), the Small Ruminant Nutrition System was used (Cannas et al., 2004). The chemical composition of orange molasses is shown in Table 2. Experimental diets were provided daily *ad libitum* as total mixed ration (TMR) to allow 10% of orts, and the average daily intake of diet was determined by daily quantify of the amount of TMR offered and of the orts throughout the experimental period. Manually the orange molasses was added to a pre-mix with the other ingredients, it was homogenized and immediately fed to the animals. The experimental period lasted 72 days, with 1 subperiod of 16 days and 2 subperiods of 28 days.

Lambs were weighed after a 16-h fast at the beginning of the experiment and on days 16, 44 and 72 of the experimental periods to determine the ADG and FE (FE; kg of BW gain/kg of DMI). At each experimental period, diets samples and orts samples were collected and frozen at -20 °C for later analysis. After the end of the trial, feed and orts samples were dried in a forced-air oven at 55 °C for 72 h (AOAC 1990; #930.15). Then all samples were ground with a Wiley mill (Marconi, Piracicaba, São Paulo, Brazil) to pass a 1mm screen. The DM was determined by oven-drying at 65 °C for 72 h and then at 105

°C for 24 h according to the method of the Association of Official Analytical Chemists (AOAC, 1990; #930.15). Ash was determined by incinerating the sample in a muffle furnace at 550 °C for 4 h (AOAC, 1990; #942.05). The total nitrogen concentration in the samples of feed and orts were determined using the Dumas combustion method using a LecoTruMac N (Leco Corporation, St. Joseph, MI USA), according to AOAC (1997; #990.03). The ether extract (EE) was determined using an Ankom XT15 extractor (Ankom Tech Corp., Macedon, NY, USA) according to AOAC (1990; #920.39). The neutral detergent fiber (NDF) concentration was determined with an Ankom A2000 Fiber Analyzer (Ankom Tech. Corp., Macedon, NY, USA; AOAC, 1990; #968.06) according to Van Soest et al. (1991). Non-fiber carbohydrates (NFC) were calculated according to equation: $NFC (g/kg) = 1000 - ((NDF (g/kg) + CP (g/kg) + EE (g/kg) + Ash (g/kg)))$.

The data were analyzed using the MIXED procedure (SAS Inst. Inc Cary, NC) and the command LSMEANS was used to generate individual means. The effects of orange molasses levels in the diets were determined by linear and quadratic polynomials. The difference was considered significant when $P < 0.05$ and trend when $P > 0.05$ and < 0.10 .

3. Results And Discussion

There was interaction between treatments and experimental periods for DMI (Table 3). In the first period the DMI decreased linearly ($P < 0.01$) and in the second period there was a trend for DMI decreased linearly ($P = 0.06$) according to the increase in orange molasses content in the diets. On the other hand, in the third period, the DMI was not affected by the treatments. It was clear that the lambs were adapting to the diets in the first two periods. However, in the third period, the similarity in DMI shows that the lambs were already completely adapted to the diets (Table 3). Probably a factor associated with the negative effect on DMI in the two initial periods was the high moisture content of orange molasses (Table 2).

There was interaction between treatments and experimental periods for ADG ($P = 0.05$). In the first period, the ADG decreased linearly ($P = 0.01$) as the orange molasses increased and in the second period they were not affected by the treatments. Otherwise, in the third period the ADG ($P = 0.05$) increased linearly as flint corn replaced orange molasses. The rate of decline in DMI when corn replaced orange molasses up to 40% was higher in the first period than in the second period (Table 3), this explains the linear decrease in ADG in the first period and similarity in ADG in the second period. It is very well documented the positive correlation between DMI and ADG (AFRC, 1998; Cannas et al., 2004; INRA, 2007; NRC, 2007; Vieira et al., 2013).

There was a trend of interaction between treatment and period for FE ($P = 0.09$). In the first period had a decreased linear effect, in the second period had no effects, on the other hand in the third period there was a trend ($P = 0.07$) of linear increased as flint corn was replaced by orange molasses. In turn, in the third period, the linear increase in ADG was consistent with the increase in FE as the content of orange molasses in the diet increased. In the present study, the overall mean ADG was similar between treatments, resulting in similar BW throughout the experiment (Table 3).

The linear increase in FE in the third period indicates that orange molasses is an ingredient with high nutritional value for lambs. Orange molasses has a high content of NFC, mainly pentoses, simple sugars, such as fructose, and soluble fiber, mainly pectin and beta-glucans (Wagner et al., 1983; Bampidis et al., 2006). These compounds are more extensively digested in the rumen when compared to the starch present in flint corn (Ariza et al., 2001), which made it possible to increase FE.

4. Conclusion

In conclusion, the orange molasses can replace up to 40% of flint corn in diets for feedlot lambs without affecting final BW. However, it is important to consider the adaptation time proved to be very important for better use of orange molasses as a source of energy in diets for lambs.

Declarations

Acknowledgments

The authors would like to thank the USP (University of São Paulo), SIPOC (Sheep and Goats Intensive Production System) for providing the facilities and financial support and CP Kelco Brazil Company financial support of this study.

Statements & Declarations

Funding

This study was funded by the FEALQ (Luiz de Queiroz Foundation), USP - ESALQ (University of São Paulo, "Luiz de Queiroz" College of Agriculture) and CP Kelco Brazil Company.

Competing Interests

We have no conflict of interest to declare.

Author contributions

Isabela J. dos Santos: Conceptualization, Methodology, Investigation, Validation, Formal analysis, Data curation, Writing original draft, **Paulo César G. Dias Junior, Rhaissa G. de Assis, Adrielly Lais Alves, Ana Carolina S. Vicente Mateus Vigo Vercesi Almada Nogueira:** Investigation, Writing – review & editing; **Janaina S. Biava and Alexandre V. Pires:** Methodology, Writing – review & editing, Visualization; **Evandro M. Ferreira:** Conceptualization, Methodology, Validation, Formal analysis, Writing – review & editing, Supervision, Funding acquisition.

Data Availability

The datasets analyzed in the current study are available from the corresponding author on a reasonable request.

Ethics approval

All the procedures have been conducted in accordance with Animal Care and Use Committee (research protocol number 7366010422).

Consent to participate

All authors consented to participation.

Consent to publish

All authors consented to submit the manuscript to the journal.

References

1. Agricultural and Food Research Council 1998. The nutrition of goats. Technical committee on response to nutrients. CAB International, Wallingford, UK
2. AOAC, 1990. Official Methods of Analysis of AOAC International, 14th ed. AOAC international, Arlington. VA.
3. AOAC, 1997. Official Methods of Analysis of AOAC International, 16th ed. AOAC international, Arlington. V.A.
4. Ariza, P., Bach, A., Stern, M.D., Hall, M.B., 2001. Effects of carbohydrates from citrus pulp and hominy feed on microbial fermentation in continuous culture 1, *J. Anim. Sci.* 79 107 - 127. <https://doi.org/10.2527/2001.79102713x>
5. Arthington, J.D., Kunkle, W.E., Martin, A.M., 2002. Citrus pulp for cattle. *Vet. Clin. North Am. - Food Anim. Pract.* 18, 317–326. [https://doi.org/10.1016/S0749-0720\(02\)00023-3](https://doi.org/10.1016/S0749-0720(02)00023-3)
6. Bampidis, V.A., Robinson, P.H., 2006. Citrus by-products as ruminant feeds: A review. *Anim. Feed Sci. Technol.* 128 175-217 <https://doi.org/10.1016/j.anifeedsci.2005.12.002>
7. Becker, R.B., Dix Arnold, P.T., Davis, G.K., Fouts, E.L., 1944. Citrus Molasses - A New Feed. *J. Dairy Sci.* 27, 269–273. [https://doi.org/10.3168/jds.S0022-0302\(44\)92594-4](https://doi.org/10.3168/jds.S0022-0302(44)92594-4)
8. Caparra, P., Foti, F., Scerra, M., Sinatra, M.C., Scerra, V., 2007. Solar-dried citrus pulp as an alternative energy source in lamb diets: Effects on growth and carcass and meat quality. *Small Rumin. Res.* 68, 303–311. <https://doi.org/10.1016/j.smallrumres.2005.11.015>
9. Cannas, A., Tedeschi, L.O., Fox, D.G., Pell, A.N., Van Soest, P.J., 2004. A mechanistic model for predicting the nutrient requirements and feed biological values for sheep. *J. Anim. Sci.* 82, 149–169. <https://doi.org/10.2527/2004.821149x>
10. Chen, M.C., Ammerman, C.B., Henry, P.R., Palmer, A.Z., Long, S.K., 1982. Citrus condensed molasses solubles as an energy source for ruminants. *J. Anim. Sci.* 53, 253–259. <https://doi.org/10.2527/jas1981.531253x>

11. De Evan, T., Cabezas, A., de la Fuente, J., Carro, M.D., 2020. Feeding agroindustrial byproducts to light lambs: Influence on growth performance, diet digestibility, nitrogen balance, ruminal fermentation, and plasma metabolites. *Animals* 10. 167 - 189 <https://doi.org/10.3390/ani10040600>
12. Ensminger, M. E., Oldfield, J. E., & Heinemann, W. W. 1990. *Feeds & nutrition: Formerly Feeds & nutrition, complete*. Clovis, Calif: Ensminger Pub. Co ISBN 0941218082 9780941218085
13. Fegeros, K., Zervas, G., Stamouli, S., Apostolaki, E., 1995. Nutritive Value of Dried Citrus Pulp and Its Effect on Milk Yield and Milk Composition of Lactating Ewes. *J. Dairy Sci.* 78, 1116–1121. [https://doi.org/10.3168/jds.S0022-0302\(95\)76728-5](https://doi.org/10.3168/jds.S0022-0302(95)76728-5)
14. Institut National de la Recherche Agronomique. 2007. *Feeding of cattle, sheep and goats*.
15. *Food Values. Tables INRA 2007*. Editions Quae, Versailles, France.
16. Mérillot, J. M., 1998. Management strategies related to legislative, economic, agronomic and environmental impact of organic wastes. *Proceeding 8th Int. Conf. Manag. Strateg. Org. Waste Use Agric.* 15–24.
17. Pires, S.R.I., de Camargo, J.B., Antonioli, P.D., 2021. A case of 55 years of servitisation in the frozen concentrated orange juice industry. *Int. J. Serv. Oper. Manag.* 39, 301–315. <https://doi.org/10.1504/IJSOM.2021.116102>
18. Santos, G.T., Lima L.S., Schogor, A.L.B., Romero, J.V., De Marchi, F.E., Grande, P.A., Santos, N.W., Santos, F.S., Kazama, R. 2014. Citrus pulp as a dietary source of antioxidants for lactating holstein cows fed highly polyunsaturated fatty acid diets. *Asian-Aust. Journal of Animal Science.*, 27, 1104–1113. <https://doi.org/10.5713/ajas.2013.13836>
19. Silvestre, A.M., Millen, D.D., 2021. The 2019 Brazilian survey on nutritional practices provided by feedlot cattle consulting nutritionists. *Rev. Bras. Zootec.* 50. 189 -201. <https://doi.org/10.37496/rbz5020200189>
20. Tuttobene, R., Avola, G., Gresta, F., Abbate, V., 2009. Industrial orange waste as organic fertilizer in durum wheat. *Agron. Sustain. Dev.* 29, 557–563. <https://doi.org/10.1051/agro/2009014>
21. USDA. 2021. National Agricultural Statistics Service: Quick stats.USDA.https://www.nass.usda.gov/Quick_Stat
22. Van Soest, P.J., Robertson, J.B., Lewis, B.A., 1991. Methods for Dietary Fiber, Neutral Detergent Fiber, and Nonstarch Polysaccharides in Relation to Animal Nutrition. *J. Dairy Sci.* 74, 3583–3597. [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
23. Vastolo, A., Calabrò, S., Cutrignelli, M.I., 2022. A review on the use of agro-industrial CO-products in animals' diets. *Ital. J. Anim. Sci.* 21, 577–594. <https://doi.org/10.1080/1828051X.2022.2039562>
24. Vieira, P.A.S., Pereira, L.G.R., Azevêdo, J.A.G., Neves, A.L.A., Chizzotti, M.L., Dos Santos, R.D., De Araújo, G.G.L., Mistura, C., Chaves, A.V., 2013. Development of mathematical models to predict dry matter intake in feedlot Santa Ines rams. *Small Rumin. Res.* 112, 78–84. <https://doi.org/10.1016/j.smallrumres.2012.10.007>
25. Wagner, J.J., Lusby, K.S., Horn, G.W., 2018. Condensed Molasses Solubles, corn steep liquor and fermented ammoniated condensed whey as protein sources for beef cattle grazing dormant native

range. J. Anim. Sci. 57, 542-552.<https://doi.org/10.2527/jas1983.573542x>

26. Wing, J.M., Van Horn, H.H., Sklare, S.D., Harris, B., 1988. Effects of citrus molasses distillers solubles and molasses on rumen parameters and lactation. J. Dairy Sci. 71, 414–420.
[https://doi.org/10.3168/jds.S0022-0302\(88\)79571-5](https://doi.org/10.3168/jds.S0022-0302(88)79571-5)

Tables

Table 1. Proportion of ingredients and chemical composition of experimental diets (g/kg of DM).

Ingredients (g/kg of DM)	Treatments ¹		
	00M	200M	400M
Hay of coastcross (<i>Cynodon</i> sp.)	100	100	100
Ground flint corn	707	565	424
Orange molasses	-	141	282
Soybean meal	143	143	142
Urea	-	1	2
Ammonium chloride	5	5	5
Limestone	14	14	14
Mineral mix ²	15	15	15
Monensin ³ , mg/kg of MS	16	16	16
Chemical Composition (g/kg of DM)			
Dry matter (g/kg as-fed basis)	912	525	415
Crude protein	166	164	164
Neutral detergent fiber	199	173	158
Ether extract	41	37	32
Non-fibrous carbohydrate	540	570	592
ME (Mcal/kg of DM) ⁴	2.9	2.7	2.8

¹00M: 0% inclusion of orange molasses in dry matter; 200M: 20% inclusion of orange molasses replacing corn; 400M: 40% inclusion of orange molasses replacing corn.

²Composition (dry matter basis): 75 g/kg P; 134 g/kg Ca; 10 g/kg Mg; 70 g/kg S; 145 g/kg Na; 500 ppm Fe; 300 ppm Cu; 4600 ppm Zn; 15 ppm Se.

³Rumensin 100 (Sodium monensin, Elanco of Brazil, São Paulo, Brazil).

⁴ME: Metabolizable energy according to *Small Ruminant Nutrition System* (Cannas et al., 2004).

Table 2. Chemical composition of orange molasses.

Chemical composition (g/kg of DM)	Orange Molasses
Dry matter (g/kg as-fed basis)	171
Crude protein	84
Neutral detergent fiber	24
Ether extract	12
Non-fibrous carbohydrate	842

Table3. Effects of increasing levels of orange molasses in replacement corn on performance of feedlot lambs.h

Item ²	Treatments ¹			SEM ³	P – Value ⁴				
	00M	200M	400M		L	Q	T	P	T × P
Body weight (kg)									
Initial, day 0	29.84	30.70	31.00	0.89	0.74	0.21			
1 st Period, day 16	32.62	32.46	31.73	1.26	0.47	0.81			
2 nd Period, day 44	39.13	39.13	37.64	1.28	0.39	0.95			
3 rd Period, day 72	44.38	44.24	44.29	1.49	0.99	0.84			
Mean	38.67	38.46	37.88	0.83	0.41	0.83	0.69	< 0.01	0.97
Dry matter intake (kg/day)									
1 st Period, day 16	0.98	0.80	0.60	0.04	< 0.01	0.94			
2 nd Period, day 44	1.15	1.03	0.96	0.07	0.06	0.81			
3 rd Period, day 72	1.10	1.17	1.18	0.06	0.17	0.90			
Mean	1.09	1.00	0.90	0.03	< 0.01	0.98	0.01	< 0.01	< 0.01
ADG (kg/day)									
1 st Period, day 16	0.19	0.12	0.03	0.04	< 0.01	0.80			
2 nd Period, day 44	0.23	0.24	0.21	0.06	0.82	0.22			
3 rd Period, day 72	0.18	0.19	0.24	0.02	0.05	0.78			
Mean	0.20	0.19	0.17	0.02	0.25	0.80	0.5	< 0.01	0.05
Feed efficiency									
1 st Period, day 16	0.19	0.14	0.05	0.04	0.08	0.41			
2 nd Period, day 44	0.19	0.21	0.21	0.02	0.69	0.80			
3 rd Period, day 72	0.16	0.16	0.20	0.02	0.07	0.10			

72									
Mean	0.18	0.18	0.19	0.01	0.64	0.15	0.64	0.02	0.09

¹OOM: 0% inclusion of orange molasses; 2OOM: 20% inclusion of orange molasses replacing corn; 4OOM: 40% inclusion of orange molasses replacing corn. ²ADG: average daily gain.

³Sem: standard error of the mean.

⁴L: linear effect; Q: quadratic effect; T: treatment effect; P: period effect; P x T: period and treatment interaction effect.